To remove capsule: 1. Remove tab. 2. Push capsule through from other side.



WARNING: SEVERE, LIFE-THREATENING HUMAN BIRTH DEFECTS. IF THALIDOMIDE IS TAKEN DURING PREGNANCY, IT CAN CAUSE BIRTH DEFECTS OR DEATH TO AN UNBORN BABY. THALIDOMIDE SHOULD NEVER BE USED BY WOMEN WHO ARE PREGNANT OR WHO COULD BECOME PREGNANT WHILE TAKING THE DRUG. EVEN A SINGLE DOSE [1 CAPSULE (50 mg)] TAKEN BY A PREGNANT WOMAN CAN CAUSE SEVERE BIRTH DEFECTS.



WARNING: THALOMID® (thalidomide) CAN CAUSE SEVERE LIFE-THREATENING HUMAN BIRTH DEFECTS.

©1999 Celgene Corporation Printed in the U.S.A. 8/99 BL105 13.001 Store at 59 to 86 °F; 15 to 30 °C. Protect from light.

General Guidelines for Taking THALOMID® (thalidomide)

DO NOT GET PREGNANT

READ ALL WARNINGS CAREFULLY BEFORE TAKING THIS MEDICATION.

Before taking THALOMID® (thalidomide), you must:

- ☐ Understand that THALOMID® (thalidomide) causes severe birth defects.
- ☐ Know how to prevent pregnancy, if you are a woman.
- ☐ Understand the need to use latex condoms during sex with a woman, if you are a man.
- ☐ Understand that you must NEVER share this drug with anyone else, even if they have symptoms like yours.
- ☐ Know about other health problems called "side effects."
- ☐ Understand that you must NEVER donate blood or sperm while you are taking this drug.
- ☐ Understand that THALOMID® (thalidomide) does not induce abortion of the fetus and should never be used for contraception.

Celgene

Manufactured for Celgene Corporation Warren, NJ 07059 USA

This package contains important information about THALOMID® (thalidomide), but does not contain all information about the drug that may be of importance to patients. If you have any questions, talk to your doctor or pharmacist.

LOT 0161S
EXP 11/02

THALOMID® (thalidomide) Each capsule contains 50 mg thalidomide. **50 mg Capsules**

50 mg 28 Capsules Each capsule contains 50 mg thalidomide. **NDC 59572-105-13**

THALOMID® (thalidomide)

Rx only

WARNING: SEVERE, LIFE-THREATENING HUMAN BIRTH DEFECTS.
IF THALIDOMIDE IS TAKEN DURING PREGNANCY, IT CAN CAUSE SEVERE BIRTH DEFECTS OR DEATH TO AN UNBORN BABY. THALIDOMIDE SHOULD NEVER BE USED BY WOMEN WHO ARE PREGNANT OR WHO COULD BECOME PREGNANT WHILE TAKING THE DRUG. EVEN A SINGLE DOSE [1 CAPSULE (50 mg)] TAKEN BY A PREGNANT WOMAN CAN CAUSE SEVERE BIRTH DEFECTS.

PHARMACIST: DISPENSE INTACT. THIS DRUG MUST NOT BE REPACKAGED. DO NOT DISPENSE UNLESS PATIENT HAS BEEN COUNSELED ON THE RISK OF BIRTH DEFECTS ASSOCIATED WITH THIS DRUG, AND A SIGNED INFORMED CONSENT FORM HAS BEEN PRESENTED.

THIS AREA FOR PRESCRIPTION LABEL

Class 5